

Dirac and Weyl Equations on a Lattice as Quantum Cellular Automata

Iwo Białynicki-Birula

Centrum Fizyki Teoretycznej PAN
Lotników 32/46, 02-668 Warsaw, Poland
and*

*Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität
Robert-Mayer-Strasse 8-10, Frankfurt am Main, Germany*

A discretized time evolution of the wave function for a Dirac particle on a cubic lattice is represented by a very simple quantum cellular automaton. In each evolution step the updated value of the wave function at a given site depends only on the values at the nearest sites, the evolution is unitary and preserves chiral symmetry. Moreover, it is shown that the relationship between Dirac particles and cellular automata operating on two component objects on a lattice is indeed very close. Every local and unitary automaton on a cubic lattice, under some natural assumptions, leads in the continuum limit to the Weyl equation. The sum over histories is evaluated and its connection with path integrals and theories of fermions on a lattice is outlined.

PACS numbers: 11.10.Qr, 11.15.Ha, 02.70.+d

The aim of this paper is to present a very simple lattice algorithm that reproduces in the continuum limit the propagation of massless or massive Dirac particles. The discretized time evolution described here satisfies the fundamental physical requirements of locality (the updated value of the wave function at a given site depends only on the values at the neighboring sites), unitarity (the norm of the wave function is preserved), and chiral symmetry (independent propagation of two helicity states).

The main, and totally unexpected, result of this investigation is a discovery that for two-component wave functions on a cubic lattice the Weyl equation necessarily follows in the continuum limit from locality, unitarity, and two additional, natural assumptions: (A) wave functions that have constant values throughout the lattice do not change in time and (B) the evolution algorithm preserves the symmetry of the lattice. Thus, the rotation group and spin emerge automatically in the continuum limit from unitary dynamics on a cubic lattice.

The results presented in this paper are directly related to numerous proposals of path integrals for a Dirac particle since an iteration of a discretized time evolution gives a sum over histories. The path integral for the Dirac particle in one space dimension was found by Feynman [1,2] and independently by Riazanov [3]. Even this relatively simple problem, where there is no spin to complicate matters, is still attracting attention [4,5]. There is no consensus at all as to the form of path integrals for a Dirac particle in three dimensions. Previously proposed path integrals fall into three broad categories. The first category [3,6–13] comprises those approaches that work by “reverse engineering” introducing from the outset the Dirac matrices to describe the spin degrees of freedom. These approaches are often combined with the proper-time representation of the Dirac operator. Into the second category [14–16] fall those formulations that derive spin from continuously parametrized space of states related to the rotation group. To the third category [17–20] belong all approaches based on anticommuting Grassmannian variables. The sum over histories that is obtained from my lattice algorithm is distinct from all these path integrals since the evolution is fully discretized and the rotation group emerges only in the continuum limit.

There is an obvious connection between this work and theories of fermions on a lattice but there is also an essential difference in the choice of objects to be calculated. I make no attempt to write down discretized versions of the Hamiltonian or the Lagrangian working instead with a discretized version of the evolution operator. In my approach common difficulties (breaking of chiral symmetry, doubling of fermion species, nonlocality) encountered in formulating the dynamics of a Dirac particle on a lattice [21–27] and the no-go theorem concerning Weyl particles on a lattice [28] are avoided.

Algorithms for cellular automata describing quantum system have been studied by Groessing and Zeilinger [29], but their discretized time evolution does not conserve probability. In a recent paper Kostin [30] has introduced a cellular automaton for the Dirac equation that does conserve probability but his algorithm is nonlinear and it gives a linear evolution equation only in the continuum limit. Following the example set by these authors I use the term *quantum cellular automaton* to denote discretized evolution of complex wave functions. I believe that the name *cellular automaton* is justified despite the fact that one of the eight properties of cellular automata required by Wolfram [31] does not hold: the states are described by continuous, not by discrete variables. However, my algorithm for updating the state of the system is synchronous, homogeneous, discrete in space and in time, deterministic, and spatially and temporally local. In my case the name *quantum* is fully justified by the unitarity of time evolution.

Let me begin with a massless particle on a cubic lattice. In my quantum cellular automaton the two-component wave function $\phi(i, j, k, t)$ is updated for each time increment Δt according to the following local algorithm

$$\begin{aligned}\phi(i, j, k, t + \Delta t) = & W_{+++}\phi(i + 1, j + 1, k + 1, t) \\ & + W_{++-}\phi(i + 1, j + 1, k - 1, t) + \cdots \\ & + W_{---}\phi(i - 1, j - 1, k - 1, t),\end{aligned}\quad (1)$$

where the integers i, j and k are the coordinates of lattice sites in units of the cell size a ($\mathbf{r} = (ai, aj, ak)$) and all eight W 's are 2×2 matrices. This algorithm can be written in a compact form

$$\phi(\mathbf{r}, t + \Delta t) = \sum_{\mathbf{h}} W(\mathbf{h})\phi(\mathbf{r} + \mathbf{h}, t), \quad (2)$$

where the summation extends over a set of vectors $\mathbf{h}$ that point from a given site to eight nearest sites. Their components in units of a are ± 1 ,

$$\begin{aligned}\{\mathbf{h}\} = & \{\{1, 1, 1\}, \{1, 1, -1\}, \{1, -1, 1\}, \{1, -1, -1\} \\ & \{-1, 1, 1\}, \{-1, 1, -1\}, \{-1, -1, 1\}, \{-1, -1, -1\}\}.\end{aligned}\quad (3)$$

Upon evaluating the norm of the updated wave function, one finds that the matrices $W(\mathbf{h})$ must satisfy the following algebraic relations to guarantee the unitarity of the transformation (2)

$$\sum_{\mathbf{h}} W^\dagger(\mathbf{h})W(\mathbf{h}) = 1, \quad (4)$$

$$\sum_{\mathbf{h}} W^\dagger(\mathbf{h})W(\mathbf{h} + \mathbf{h}' - \mathbf{h}'') = 0, \quad (5)$$

where $\mathbf{h}'$ and $\mathbf{h}'' \neq \mathbf{h}'$ are arbitrary vectors belonging to the set (3) and the sum in (5) extends only over vectors $\mathbf{h}$ such that the vectors $\mathbf{h} + \mathbf{h}' - \mathbf{h}''$ are also members of the set. It follows from these unitarity conditions that the inverse of (2) has a similar local form so that my automaton is fully reversible,

$$\phi(\mathbf{r}, t) = \sum_{\mathbf{h}} W^\dagger(\mathbf{h})\phi(\mathbf{r} + \mathbf{h}, t + \Delta t). \quad (6)$$

Since the inverse of a unitary transformation is also unitary, the hermitian conjugate matrices $W^\dagger$ must obey the same conditions as the matrices W ,

$$\sum_{\mathbf{h}} W(\mathbf{h})W^\dagger(\mathbf{h}) = 1, \quad (7)$$

$$\sum_{\mathbf{h}} W(\mathbf{h})W^\dagger(\mathbf{h} + \mathbf{h}' - \mathbf{h}'') = 0. \quad (8)$$

Depending on the mutual orientation of the vectors $\mathbf{h}'$ and $\mathbf{h}''$, the conditions (5) and (8) have one, two, or four terms, as exemplified below

$$W_{+++}^\dagger W_{---} = 0, \quad (9a)$$

$$W_{+++}^\dagger W_{+--} + W_{-++}^\dagger W_{---} = 0, \quad (9b)$$

$$\begin{aligned}W_{+++}^\dagger W_{++-} + W_{+-+}^\dagger W_{+--} + W_{-++}^\dagger W_{-+-} \\ + W_{--+}^\dagger W_{---} = 0.\end{aligned}\quad (9c)$$

In total there are 8 conditions of the type (9a), 12 conditions of the type (9b), 6 conditions of the type (9c), and one condition (4) and then the same number of conditions with W and $W^\dagger$ interchanged. Nonetheless they can all be satisfied by the following simple and symmetric choice of W 's

$$\begin{aligned}W_{+++} &= q_+ P_1, \quad W_{++-} = q_- P_2, \quad W_{+-+} = q_- P_1, \\ W_{+--} &= q_+ P_2, \quad W_{-++} = q_- P_3, \quad W_{-+-} = q_+ P_4, \\ W_{--+} &= q_+ P_3, \quad W_{---} = q_- P_4,\end{aligned}\quad (10)$$

where

$$q_+ = (1+i)/4, \quad q_- = (1-i)/4, \quad (11)$$

and

$$\begin{aligned} P_1 &= \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}, \quad P_2 = \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}, \\ P_3 &= \begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix}, \quad P_4 = \begin{pmatrix} 0 & -1 \\ 0 & 1 \end{pmatrix}. \end{aligned} \quad (12)$$

In order to find the continuum limit of the evolution equation I shall use the exponential representation of the shift operators to write (2) in the form

$$\phi(\mathbf{r}, t + \Delta t) = \sum_{\mathbf{h}} W(\mathbf{h}) \exp(\mathbf{h} \cdot \nabla) \phi(\mathbf{r}, t). \quad (13)$$

Expanding both sides of this equation in powers of Δt and a ($\mathbf{h}$ is of the order of a) and keeping only linear terms, I obtain the Weyl equation,

$$\partial_t \phi(\mathbf{r}, t) = c \boldsymbol{\sigma} \cdot \nabla \phi(\mathbf{r}, t), \quad (14)$$

where $c = a/\Delta t$. Incidentally, to obtain a discretized Weyl equation in two space dimensions one may choose the four matrices W_{++}, W_{+-}, W_{-+} , and W_{--} as equal to the matrices P_i .

I shall show now that *for every set* of 2×2 matrices satisfying the unitarity conditions, and not just for the particular choice (10), one obtains the Weyl equation in the continuum limit. To prove this let me first expand the exponential operator in (13) into powers of a ,

$$\sum_{\mathbf{h}} W(\mathbf{h}) \exp(\mathbf{h} \cdot \nabla) = 1 + a \mathbf{s} \cdot \nabla + \dots, \quad (15)$$

where I have made use of the assumption A that a homogeneous wave function does not change in time,

$$\sum_{\mathbf{h}} W(\mathbf{h}) = 1, \quad (16)$$

and I have introduced the matrices s_i ,

$$\begin{aligned} s_x &= W_{+++} + W_{++-} + W_{+-+} + W_{+--} \\ &\quad - W_{-++} - W_{-+-} - W_{--+} - W_{---}, \end{aligned} \quad (17a)$$

$$\begin{aligned} s_y &= W_{+++} + W_{++-} - W_{+-+} - W_{+--} \\ &\quad + W_{-++} + W_{-+-} - W_{--+} - W_{---}, \end{aligned} \quad (17b)$$

$$\begin{aligned} s_z &= W_{+++} - W_{++-} + W_{+-+} - W_{+--} \\ &\quad + W_{-++} - W_{-+-} + W_{--+} - W_{---}. \end{aligned} \quad (17c)$$

Since the differentiations are antihermitian operators, the unitarity of the operator (15) requires that the three matrices s_i must be hermitian and, therefore, the formulas (17) must also hold with all the matrices W replaced by their hermitian conjugates. This fact will enable me to use the unitarity conditions (5) and (8) to calculate products of the matrices s_i , ($i = x, y, z$). The unitarity conditions also imply (to see this, one has to write them down explicitly) that all products of a W matrix by its hermitian conjugate commute, and therefore they can be simultaneously diagonalized. From the equivalence of all six lattice directions (assumption B) I conclude that the eigenvalues of all matrices $W^\dagger(\mathbf{h})W(\mathbf{h})$ must be the same and then from equations (4) and (7) I find that these eigenvalues are $1/4$ and 0 . It is now a matter of tedious but otherwise straightforward algebraic manipulations to show that the matrices s_i satisfy the familiar anticommutation relations,

$$s_i s_j + s_j s_i = 2\delta_{ij}. \quad (18)$$

There exist only two inequivalent two-dimensional representations of these relations: $\mathbf{s} = \boldsymbol{\sigma}$ or $\mathbf{s} = -\boldsymbol{\sigma}$. They describe the propagation of two helicities. Thus, up to a choice of helicity, the universality of the Weyl equation under the listed assumptions is established.

For massless particles each helicity state propagates independently. Chiral invariance (or CP symmetry) is expressed in my discretized form of time evolution by the fact that the matrices W corresponding to the two helicities are related by the spatial reflection

$$W(\mathbf{h}) \rightarrow W(-\mathbf{h}), \quad (19)$$

as seen from the formulas (17). For massive particles, however, the two helicity states are mixed by the mass term. Therefore, the discretized time evolution for a massive particle must be described, as in the standard Dirac equation, in terms of two two-component wave functions. The discrete time-evolution algorithm for a massive Dirac particle can again be written in the same general form (6) as for the massless case

$$\psi(\mathbf{r}, t + \Delta t) = \sum_{\mathbf{h}} D(\mathbf{h}) \psi(\mathbf{r} + \mathbf{h}, t). \quad (20)$$

The 4×4 matrices D that act on four-component wave functions $\psi(\mathbf{r}, t)$ can be expressed in terms of the matrices $W(\mathbf{h})$ and $W(-\mathbf{h})$ by the following block-matrix formulas (from now on I set $c = \Delta t/a = 1$ and $\hbar = 1$)

$$D(\mathbf{h}) = \begin{pmatrix} \cos(ma) & i \sin(ma) \\ i \sin(ma) & \cos(ma) \end{pmatrix} \begin{pmatrix} W(\mathbf{h}) & 0 \\ 0 & W(-\mathbf{h}) \end{pmatrix}. \quad (21)$$

One may check that such a modification does not affect the unitarity conditions; the matrices D will also satisfy all of them. The continuum limit of (20) gives the Dirac equation in the Weyl representation of the Dirac matrices

$$\partial_t \psi(\mathbf{r}, t) = (\rho_3 \boldsymbol{\sigma} \cdot \nabla + i \rho_1 m) \psi(\mathbf{r}, t). \quad (22)$$

I shall now write down the sum over histories for the Weyl particle arising from my time-evolution algorithm (2) and I shall directly show that it yields the propagator in the continuum limit. The N -fold iteration of the single-step evolution leads to the formula

$$\begin{aligned} & \phi(\mathbf{r}, t + N\Delta t) \\ &= \sum_{\mathbf{h}_1, \dots, \mathbf{h}_N} W(\mathbf{h}_1) \cdots W(\mathbf{h}_N) \phi(\mathbf{r} + \mathbf{h}_1 + \cdots + \mathbf{h}_N, t). \end{aligned} \quad (23)$$

When the sum in this formula is restricted to only those combinations of the vectors $\mathbf{h}_i$ that produce a given displacement, we will obtain a discrete version of the propagator

$$K(\mathbf{r} - \mathbf{r}', t) = \sum_{\mathbf{h}_1, \dots, \mathbf{h}_N} W(\mathbf{h}_1) \cdots W(\mathbf{h}_N) \delta_{\mathbf{r}, \mathbf{r}' + \mathbf{h}_1 + \cdots + \mathbf{h}_N}, \quad (24)$$

where $t = N\Delta t$. Each term in this sum corresponds to a lattice trajectory that in N steps connects the initial and the final lattice sites. A single step described by a vector $\mathbf{h}$ is represented by the matrix $W(\mathbf{h})$. Contributions from all trajectories are coherently added.

The constraint on the sum in (24) can be handled (cf., for example, [32]) with the help of the Fourier representation of the Kronecker delta, leading to the following integral form of the propagator

$$K(\mathbf{r} - \mathbf{r}', t) = \left(\frac{a}{2\pi}\right)^3 \int d^3 k \tilde{W}^{\frac{t}{\Delta t}}(\mathbf{k}) \exp[i\mathbf{k} \cdot (\mathbf{r} - \mathbf{r}')], \quad (25)$$

where

$$\tilde{W}(\mathbf{k}) = \sum_{\mathbf{h}} W(\mathbf{h}) \exp(i\mathbf{k} \cdot \mathbf{h}), \quad (26)$$

and the integration extends over the Brillouin zone: $-\pi/a < k_i < \pi/a$.

The integral (25) can be evaluated with the use of the explicit representation (10) of the matrices W ,

$$\begin{aligned} \tilde{W}(\mathbf{k}) &= c_x c_y c_z + s_x s_y s_z + i[(s_x c_y c_z - c_x s_y s_z) \sigma_x \\ &\quad + (c_x s_y c_z + s_x c_y s_z) \sigma_y + (c_x c_y s_z - s_x s_y c_z) \sigma_z], \end{aligned} \quad (27)$$

where

$$c_i = \cos(k_i a), \quad s_i = \sin(k_i a). \quad (28)$$

This is a unitary matrix and its eigenvalues $\lambda_{\pm}$ are

$$\begin{aligned} \lambda_{\pm} &= \exp(\pm i\varphi) \\ &= c_x c_y c_z + s_x s_y s_z \pm i \sqrt{1 - (c_x c_y c_z + s_x s_y s_z)^2}. \end{aligned} \quad (29)$$

An easy way to evaluate the N -th power ($N = t/\Delta t$) of the matrix (27), needed to calculate the propagator, is to write $\tilde{W}$ in terms of its eigenvalues and its unitary diagonalizing matrix V ,

$$\tilde{W}(\mathbf{k}) = V \begin{pmatrix} \exp(i\varphi) & 0 \\ 0 & \exp(-i\varphi) \end{pmatrix} V^\dagger, \quad (30)$$

$$(\tilde{W}(\mathbf{k}))^{\frac{t}{\Delta t}} = V \begin{pmatrix} \exp(it\varphi/a) & 0 \\ 0 & \exp(-it\varphi/a) \end{pmatrix} V^\dagger. \quad (31)$$

In the limit, when $a \rightarrow 0$, one obtains

$$\lim_{a \rightarrow 0} (\varphi/a) = |\mathbf{k}| =: k, \quad (32)$$

$$\lim_{a \rightarrow 0} V = (\mathbf{k} \cdot \boldsymbol{\sigma} - k\sigma_z) / \sqrt{2k(k - k_z)}, \quad (33)$$

and finally

$$\lim_{\Delta t \rightarrow 0} (\tilde{W}(\mathbf{k}))^{\frac{t}{\Delta t}} = \exp(i\mathbf{k} \cdot \boldsymbol{\sigma} t), \quad (34)$$

in accordance with the Weyl equation. Thus, the sum over histories (23) reproduces correctly the propagator in the continuum limit.

Having reproduced the propagation of a massless particle I can easily include the mass in the sum over histories since that part has been already solved by Feynman [1,2] in one dimension. More recently Feynman's "checkerboard" picture of a particle zigzagging through spacetime, reversing its helicity at each bend, has also been described by a Poisson process [9,15,20,32]. This Poisson process must be combined with the propagation of definite helicity states. Therefore, the propagator for a massive Dirac particle will be a sum of terms, each term describing a fixed number of helicity reversals. Between the reversals the propagation is described by the sum over histories (24) separately for each helicity.

Interaction with the electromagnetic field may be accounted for in the standard fashion by changing the phase of the wave function at each propagation step according to the local value of the electromagnetic potential.

I would like to thank Jerzy Kijowski and Joachim Reinhardt for discussions and Walter Greiner for his hospitality at the University of Frankfurt.

* Permanent address.

- [1] R. P. Feynman, Rev. Mod. Phys. **20**, 267 (1948). Unpublished notes by Feynman on path integrals for the Dirac equation in one dimension are described in S. S. Schweber, Rev. Mod. Phys. **58**, 449 (1986).
- [2] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, (McGraw-Hill, New York, 1965).
- [3] G. V. Riazanov, Zh. Eksp. Theor. Fiz. **33**, 1437 (1957) [Sov. Phys. J.E.T.P. **6**, 1107 (1958)].
- [4] T. Nakamura, J. Math. Phys. **32**, 457 (1991).
- [5] D. G. C. McKeon and G. N. Ord, Phys. Rev. Lett. **69**, 3 (1992).
- [6] G. J. Papadopoulos and J. T. Devreese, Phys. Rev. D **13**, 2227 (1976).
- [7] G. Rosen, Phys. Rev. A **28**, 1139 (1983).
- [8] M. A. Kaye and A. Inomata, Phys. Rev. Lett. **53**, 107 (1984).
- [9] B. Gaveau, T. Jacobson, M. Kac, and L. S. Schulman, Phys. Rev. Lett. **53**, 419 (1984).
- [10] A. O. Barut and I. H. Duru, Phys. Rev. Lett. **53**, 2355 (1984).

- [11] T. Ichinose and H. Tamura, J. Math. Phys. **25**, 1810 (1984).
- [12] A. M. Polyakov, in *Fields, Strings, and Critical Phenomena*, Les Houches XLIX, edited by E. Brézin and J. Zinn-Justin, (North-Holland, Amsterdam, 1990).
- [13] J. Ambjørn, B. Durhuus, and T. Jonsson, Nucl. Phys. B **330**, 509 (1989).
- [14] L. S. Schulman, Phys. Rev. **176**, 1558 (1968).
- [15] T. Jacobson, J. Phys. A **17**, 2433 (1984).
- [16] T. Jacobson, Phys. Lett. B **216**, 150 (1989).
- [17] F. A. Berezin and M. S. Marinov, Ann. Phys. (NY) **104**, 332 (1977).
- [18] F. Ravndal, Phys. Rev. D **21**, 2823 (1980).
- [19] A. T. Ogielski and J. Sobczyk, J. Math. Phys. **22**, 2060 (1981).
- [20] B. Gaveau and L. S. Schulman, Phys. Rev. D **36**, 1135 (1987).
- [21] K. G. Wilson, Phys. Rev. D **10**, 2445 (1974).
- [22] J. B. Kogut and L. Susskind, Phys. Rev. D **11**, 395 (1975).
- [23] L. Susskind, Phys. Rev. D **16**, 3031 (1977).
- [24] S. D. Drell, M. Weinstein, and S. Yankielowicz, Phys. Rev. D **14**, 487, 1627 (1976).
- [25] J. M. Rabin, Phys. Rev. D **24**, 3218 (1981).
- [26] C. M. Bender, K. A. Milton, and D. S. Sharp, Phys. Rev. D **31**, 383 (1985).
- [27] J. Kijowski and A. Thielmann, in *Differential Geometry, Group Representations, and Quantization*, edited by J. P. Hennig *et al.*, (Springer, Berlin, 1991).
- [28] H. B. Nielsen and M. Ninomiya, Nucl. Phys. B **185**, 20 (1981). The authors state explicitly in this paper that one possibility to avoid their no-go theorem is to introduce discretized time.
- [29] G. Groessing and A. Zeilinger Physica D **31**, 70 (1988).
- [30] M. D. Kostin, J. Phys. A **26**, L209 (1993).
- [31] S. Wolfram, *Theory and Applications of Cellular Automata* (World Scientific, Singapore, 1986), p.1.
- [32] T. Jacobson and L. S. Schulman, J. Phys. A **17**, 375 (1984).